



GUIDELINE

Pulmonary Haemorrhage

Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

Child Safe Organisation Statement of Commitment

CAHS commits to being a child safe organisation by applying the National Principles for Child Safe Organisations. This is a commitment to a strong culture supported by robust policies and procedures to reduce the likelihood of harm to children and young people.

This document should be read in conjunction with this [disclaimer](#)

Background

This is better termed “haemorrhagic pulmonary oedema” and is dependent on the primary cause. It is a form of fulminant lung oedema with leakage of red cells and capillary filtrate into the lungs. It must be differentiated from occurrence of a small amount of blood being aspirated from the end of an ETT secondary to trauma. It represents the extreme end of the spectrum of pulmonary oedema in the neonate.

Associated Clinical Conditions

- Severe birth asphyxia.
- Rhesus haemolytic disease with hydrops/near hydrops.
- Left heart failure.
- Congenital heart disease.
- Sepsis.
- Hypothermia.
- Fluid overload.
- Oxygen toxicity.
- Haemostasis failure.
- Any pre-existing lung disease will worsen, as protein-rich fluid in the alveoli will inhibit surfactant function.

Clinical Presentation

The commonest clinical presentation is in an infant with severe RDS on IPPV in high oxygen and heart failure secondary to a large pulmonary blood flow from a PDA. It is also associated with surfactant therapy.

Symptoms

- Sudden deterioration.
- Copious bloody secretions from the infant's airway either up the ETT or from the larynx and mouth if not already intubated.
- Usually there is hypotension.
- The infant may be pale and unresponsive.
- The outcome is dependent on the cause of the oedema. At the time of collapse, infants are susceptible to neurological damage and GMH/IVH. The mortality is significant, can be as high as 50% unless managed well.



Investigations

- Check Hb, coagulation screen.
- Check ABG, biochemistry.
- CXR: often shows a white-out.
- Look for sepsis.

Management

- Maintain blood pressure.
- Correct acidosis by ventilation or drug therapy.
- Fluid balance - consider fluid restriction, administer **Frusemide** 1-1.5 mg/kg.
- Sedate as needed.
- Use high PEEP - 6-7 cms (redistributes lung water back into the interstitial space, improving oxygenation and ventilation-perfusion balance). Consider alternative modes of ventilation; HFOV or HFJV which may enable the use of higher PEEP or MAP.
- Surfactant - may help to overcome protein inhibition of endogenous surfactant.
- Check for PDA - opening of symptomatic PDA. Consider medical treatment for PDA if neonate not requiring inhaled nitric oxide.
- Suction - be careful as may aggravate bleed.
- Manage coagulopathy.
- Sepsis - use or review antibiotics.

This document can be made available in alternative formats on request.

Document Owner:	Neonatology		
Reviewer / Team:	Neonatology Coordinating Group		
Date First Issued:	June 2006	Last Reviewed:	June 2022
Amendment Dates:		Next Review Date:	June 2023
Approved by:	Neonatology Coordinating Group	Date:	28 th June 2022
Endorsed by:	Neonatology Coordinating Group	Date:	
Standards Applicable:	NSQHS Standards:   Child Safe Standards: 1,10		

Printed or personally saved electronic copies of this document are considered uncontrolled



Healthy kids, healthy communities

Compassion Excellence Collaboration Accountability Equity Respect

Neonatology | Community Health | Mental Health | Perth Children's Hospital